

Cosmological dynamics of Scalar–Tensor Gravity

S Carloni [†], J A Leach [†], S Capozziello [‡], P K S Dunsby ^{†§}

[†] Department of Mathematics and Applied Mathematics, University of Cape Town, Rondebosch, 7701, South Africa

[‡] Dipartimento di Scienze Fisiche and INFN Sez. di Napoli, Università di Napoli ‘Federico II’, Complesso Universitario di Monte S Angelo, Via Cinthia I-80126 Napoli, Italy

[§] South African Astronomical Observatory, Observatory, Cape Town, South Africa

E-mail: sante.carloni@gmail.com, capozziello@na.infn.it, leachj@maths.uct.ac.za and pksd@maths.uct.ac.za

Abstract. We study the phase-space of FLRW models derived from Scalar–Tensor Gravity where the non-minimal coupling is $F(\phi) = \xi\phi^2$ and the effective potential is $V(\phi) = \lambda\phi^n$. Our analysis allows to unfold many feature of the cosmology of this class of theories. For example, the evolution mechanism towards states indistinguishable from GR is recovered and proved to depend critically on the form of the potential $V(\phi)$. Also, transient almost–Friedmann phases evolving towards accelerated expansion and unstable inflationary phases evolving towards stable ones are found. Some of our results are shown to hold also for the String-Dilaton action.

PACS numbers: 98.80.JK, 04.50.+h, 05.45.-a

1. Introduction

Scalar–Tensor Gravity (STG) has become a major area of activity in the last thirty years. This is due mainly to the discovery of several shortcomings of General Relativity (GR) in cosmology and quantum field theory. In fact, the presence of a big bang singularity, the flatness and horizon problems, led to the realization that the “standard” model for cosmology [1], based on GR and the standard model of particle physics, is not able to describe the universe in extreme regimes. On the other hand, GR is a *classical* theory which has so far eluded every attempt to be quantized, making the description of gravity on microscopic scales very difficult. This has led many authors to believe that the standard Einstein scheme is inadequate to fully describe the gravitational interaction at a fundamental level [2] and has stimulated the proposal and the study of alternative theories of gravity that might be able to solve these problems and at the same time reduce to GR in the weak field regime.

One of the most fruitful theories of this type is STG, in which scalar field(s) that are non-minimally coupled to gravity are introduced. These type of theories were proposed nearly half a century ago by Jordan [3] and later refined by Brans and Dicke [4]. The original motivation behind Brans-Dicke theory (BD) came from Mach’s principle, but over the years, BD and STG have gained interest in a wide number of scenarios. For example, in unification schemes such as superstrings, supergravity or

grand unified theories, the one-loop or higher-loop corrections in the high-curvature regime take the form of non-minimal couplings to the geometry or higher-order curvature invariants in their low energy effective actions [6].

In cosmology, STG acquired considerable interest because they introduce naturally scalar fields and scalar fields which are capable, among other things, of giving rise to inflationary behaviour [7]. In GR the introduction of this type of field has the drawback of raising the issue of explaining its origin. Instead, in STG this problem finds a natural solution, because the scalar field can be considered an additional degree of freedom of the gravitational interaction. For these reasons, among others, inflationary models based on STG have been widely studied [8,9].

Recently, STG has also been used to model dark energy, because scalar fields are also the natural candidates for phantom and quintessence fields (see [10] for a review). This suggests that both inflation and dark energy could be the result of the action of the same scalar field. STG offers, in this sense, the ideal framework to implement this intriguing idea and might even allow us to overcome the problems due to the energy scale difference between the dark energy scalar fields and the inflaton [11].

In this paper, we consider the problem of determining the global dynamics of Friedmann–Lemaître–Robertson–Walker (FLRW) cosmologies of STG. We will study a generic class of STG theories, where quadratic non-minimal couplings to gravity and self-interaction power-law potentials are assumed. This class of models is strictly related to the String Dilaton action and naturally exhibit duality in the cosmological solutions [12]. Furthermore, they can be obtained from generic non-minimally coupled scalar–tensor Lagrangians if Noether symmetries are found in the dynamics [13,14]. Several exact solutions of these models have been found, but the stability and global behaviour is still not well understood. Our aim is to give a full description of the global dynamics of this class of STG and determine if cosmic histories are possible that (i) present a transient matter-dominated Friedmann phase and then evolve towards an accelerated (dark energy-dominated or Λ CDM) regime or (ii) present a first unstable inflationary phase and a second inflationary attractor.

The dynamical systems approach has been already used with great success in BD [16–20], in specific models of STG [21,26–30] and in other contexts [31–33] allowing the derivation of new exact solutions and making possible a qualitative global description of the dynamics. The geometric structure and dimensionality of the phase-space of general couplings and potentials has recently been investigated by Faraoni [34].

The outline of this paper is as follows. In section 2, we derive the field equations and discuss the non-minimal coupling and the self-interacting potential. The cosmological equations for a FLRW metric are derived in section 3. We convert them into an autonomous dynamical system in order to pursue a phase-space investigation. Sections 4 and 5 are respectively devoted to the analysis of this system in the absence and presence of a perfect fluid source. We study the stability of the fixed points with the aim of identifying physically interesting cosmological behaviours. In section 6, the results are discussed and some conclusions are drawn.

The following conventions are adopted: the metric signature is $(+ - - -)$; Latin indices run from 0 to 3; ∇ is the covariant derivative; physical units $c = 8\pi G = 1$ are used.

2. The field equations

In four dimensions, a general action in which gravity is non-minimally coupled to a scalar field ϕ , reads [14]:

$$\mathcal{A} = \int dx^4 \sqrt{-g} \left[F(\phi) R + \frac{1}{2} g^{ab} \nabla_a \phi \nabla_b \phi - V(\phi) + \mathcal{L}_M \right], \quad (1)$$

where $F(\phi)$ is a generic coupling, $V(\phi)$ is the self-interaction potential and $\mathcal{L}_M$ is the matter Lagrangian.

By varying the action (1) with respect to the metric g_{ab} , we obtain the field equations

$$\begin{aligned} F(\phi) R_{ab} - \frac{1}{2} F(\phi) R g_{ab} + \frac{1}{2} \nabla_a \phi \nabla_b \phi - \frac{1}{4} g_{ab} \nabla^c \phi \nabla_c \phi \\ + \frac{1}{2} g_{ab} V(\phi) - \nabla_a \nabla_b F(\phi) + g_{ab} \square F(\phi) = -T_{ab}^M \end{aligned} \quad (2)$$

and the variation with respect to ϕ gives the Klein-Gordon equation

$$\square \phi - R F'(\phi) + V'(\phi) = 0, \quad (3)$$

where primes denote differentiation with respect to ϕ . The field equations (2) can be recast in the standard form

$$G_{ab} = R_{ab} - \frac{1}{2} R g_{ab} = T_{ab}^{eff},$$

where the effective stress-energy momentum tensor T_{ab}^{eff} is given by

$$\begin{aligned} T_{ab}^{eff} = F(\phi)^{-1} \left[-\frac{1}{2} \nabla_a \phi \nabla_b \phi + \frac{1}{4} g_{ab} \nabla^c \phi \nabla_c \phi - \frac{1}{2} g_{ab} V(\phi) \right. \\ \left. + \nabla_a \nabla_b F(\phi) - g_{ab} \square F(\phi) - T_{ab}^M \right]. \end{aligned} \quad (4)$$

This form of the equations shows clearly one of the most interesting features of Scalar Tensor Gravity: the Newtonian gravitational constant G_N , intended as the coupling of gravity with standard matter, has to be replaced by an effective gravitational "constant" G_{eff} that depends on the non-minimal coupling F (and, as a consequence, on ϕ) and *varies* in time.

The Bianchi identities $G^a{}_{b;a} = 0$ give the conservation laws for both the matter and the scalar field. As a general result [14], it is possible to show that the conservation law for the scalar field is the Klein-Gordon equation.

The action (1) is very general and contains several interesting physical cases. For example, considering the transformation

$$\phi \rightarrow \exp[-\psi], \quad F(\phi) \rightarrow \frac{1}{2} \exp[-2\psi], \quad V(\phi) \rightarrow \frac{1}{2} \Lambda \exp[-2\psi], \quad (5)$$

which specifies a particular form of coupling and potential, leads to the 4D-String-Dilaton Effective Action

$$\mathcal{A} = \int dx^4 \sqrt{-g} e^{-\psi} \left[R + g^{ab} \nabla_a \psi \nabla_b \psi - \Lambda \right], \quad (6)$$

where Λ is the string charge. In this context, such a theory is nothing else but a particular STG [35]. This means that the considerations and results for the action (1), and the related dynamics, also hold for string-dilaton cosmology. On the other hand, the set of transformations

$$F(\phi) \rightarrow \psi, \quad \frac{F(\phi)}{2F'(\phi)^2} \rightarrow -\omega(\psi), \quad V(\phi) \rightarrow 0, \quad (7)$$

give rise to the action

$$\mathcal{A} = \int dx^4 \sqrt{-g} \left[\psi R - \frac{\omega(\psi)}{\psi} g^{ab} \nabla_a \psi \nabla_b \psi \right], \quad (8)$$

which is nothing else but the BD action[‡] [4]. In addition to the ones above other interesting kind of transformations are possible [5].

In this paper, we will focus on models where the non-minimal coupling has the form $F(\phi) = \xi \phi^2$ and the self-interaction potential is taken to be an arbitrary power law of the form $V(\phi) = \lambda \phi^n$. This choice is general and motivated by several mathematical and physical reasons. In particular, beside the string-dilaton and BD actions, several effective quantum field theories, in low energy physics, can be related to such couplings and self-interacting potentials [2, 36]. Furthermore, this coupling and potential satisfy the requirement of Noether symmetries for the Lagrangian in (1), giving rise to general exact solutions of physical interest [13].

A final remark concerns the parameters of the theory. Very different models can be parameterized by the set (ξ, λ, n) , but not all the combinations of these parameters are necessarily physical. For example, attractive gravity is achieved for $\xi < 0$ and $\lambda > 0$, although physically interesting situation can also be achieved for $\xi > 0$, $\lambda < 0$ [37]. In this paper, we will consider only the physical case $\xi < 0$, $\lambda > 0$. The discussion of the other interesting combinations of parameters can be found in [38].

3. The FLRW dynamical system

In order to analyze the phase-space of the Scalar Tensor FLRW cosmologies, the field equations need to be recast in a dynamical system form. In the FLRW metric and with our choice of coupling and potential the the above Einstein-Klein-Gordon equations reduce to:

$$\frac{\ddot{a}}{a} + H \frac{\dot{\phi}}{\phi} + \frac{\ddot{\phi}}{\phi} + \left(1 - \frac{1}{6\xi}\right) \frac{\dot{\phi}^2}{\phi^2} + \frac{\lambda}{6\xi} \phi^{n-2} - \frac{\mu}{6\xi \phi^2} (1 + 3w) = 0, \quad (9)$$

$$H^2 + \frac{k}{a^2} + 2H \frac{\dot{\phi}}{\phi} + \frac{\lambda}{6\xi} \phi^{n-2} + \frac{1}{12\xi} \frac{\dot{\phi}^2}{\phi^2} + \frac{\mu}{3\xi \phi^2} = 0, \quad (10)$$

$$\frac{\ddot{\phi}}{\phi} + 3H \frac{\dot{\phi}}{\phi} + 12\xi \left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) + n\lambda \phi^{n-2} = 0, \quad (11)$$

where the first is the cosmological equation for the acceleration, the second is the Hamiltonian constraint, i.e. the $\{0,0\}$ equation, and the third is the Klein-Gordon equation. $H = \dot{a}/a$ is the Hubble parameter and k is the spatial curvature constant. We have also assumed standard matter to be a perfect fluid with a barotropic index w , so that the conservation equation yields

$$\dot{\mu} = -3H\mu(1+w), \quad (12)$$

where μ is the the matter-energy density. In what follows, since we want to stress the role of a non minimally coupled scalar field in the modelling of dark energy, we will consider only $0 < w < 1$. Other values of these parameters, which can be associated with more exotic forms of matter energy densities, although interesting, will not be considered here.

[‡] To be precise, the proper BD action is exactly recovered only for $\omega = \text{constant}$.

The equations above can be converted into an autonomous system of first-order differential equations by defining the following set of expansion normalised variables:

$$\begin{aligned} x &= \frac{\dot{\phi}}{\phi H}, & y &= \frac{\lambda \phi^{n-2}}{6\xi H^2}, \\ z &= \frac{\mu}{3\xi \phi^2 H^2}, & K &= \frac{k}{a^2 H^2}. \end{aligned} \quad (13)$$

from which we obtain

$$\begin{aligned} x' &= \frac{1}{12\xi - 1} \left[12\xi(1 + K + Kx) + 2(1 + 6\xi)x + (5 - 24\xi)x^2 - \left(1 - \frac{1}{6\xi}\right)x^3 \right. \\ &\quad \left. + 6\xi(n - 2)y - (1 - 6n\xi)xy + 6\xi(1 + 3w)z + \frac{1}{2}(1 + 3w)xz \right], \\ y' &= \frac{2y}{12\xi - 1} [24\xi - 1 + 12\xi K + 4x + (12\xi - 1)(n - 2)x \\ &\quad - \left(1 - \frac{1}{6\xi}\right)x^2 - (1 - 6n\xi)y + \frac{1}{2}(1 + 3w)z], \\ z' &= \frac{z}{12\xi - 1} [7(1 - 12\xi) - 2 - 3w(1 - 12\xi) + 24\xi K + 6(1 - 4\xi)x \\ &\quad - 2\left(1 - \frac{1}{6\xi}\right)x^2 - 2(1 - 6n\xi)y + (1 + 3w)z], \\ K' &= \frac{2K}{12\xi - 1} \left[12\xi(1 + K) + 2x - \left(1 - \frac{1}{6\xi}\right)x^2 - (1 - 6n\xi)y + \frac{1}{2}(1 + 3w)z \right], \end{aligned} \quad (14)$$

where primes denote derivatives with respect to a new evolution variable $\tau = \ln a$ and the dynamical variables are constrained by

$$1 + 2x + \frac{1}{12\xi}x^2 + y + K - z = 0. \quad (15)$$

The associated phase-space is 4-dimensional and the evolution is constrained by (15). The task is now to study the structure of such a space: this means finding the stability of the fixed points, and then to analyze the evolution of trajectories [15]. We will consider two cases: the vacuum case ($\mu = 0$) and the matter case ($\mu \neq 0$).

4. The vacuum case

When we consider the vacuum case ($\mu = 0$), the set of dynamical equations (14) reduces to

$$\begin{aligned} x' &= \frac{1}{12\xi - 1} \left[12\xi(1 + K + Kx) + 2(1 + 6\xi)x + (5 - 24\xi)x^2 - \left(1 - \frac{1}{6\xi}\right)x^3 \right. \\ &\quad \left. + 6\xi(n - 2)y - (1 - 6n\xi)xy \right], \\ y' &= \frac{2y}{12\xi - 1} [24\xi - 1 + 12\xi K + 4x + (12\xi - 1)(n - 2)x \\ &\quad - \left(1 - \frac{1}{6\xi}\right)x^2 + (6n\xi - 1)y], \\ K' &= \frac{2K}{12\xi - 1} \left[12\xi(1 + K) + 2x - \left(1 - \frac{1}{6\xi}\right)x^2 + (6n\xi - 1)y \right], \end{aligned} \quad (16)$$

with the constraint equation given by

$$1 + 2x + \frac{1}{12\xi}x^2 + y + K = 0. \quad (17)$$

In this case, the phase-space is 3-dimensional.

4.1. Finite analysis

We can further simplify the system (16) by implementing the constraint (17):

$$\begin{aligned} x' &= -2x - 4x^2 - \frac{1}{6\xi}x^3 + \frac{y}{12\xi - 1} [6\xi(n-4) + (6\xi(n-2) - 1)x], \\ y' &= y \left[2 + (n-6)x - \frac{1}{3\xi}x^2 + \frac{2(6\xi(n-2) - 1)}{12\xi - 1}y \right]. \end{aligned} \quad (18)$$

From these equations, it is clear that the x axis ($y = 0$), characterized by the absence of a potential for the scalar field, is an invariant submanifold. This tells us that there is no orbit for which the potential of the scalar field can become exactly zero and that if the potential is initially set to zero, it will remain zero.

The fixed points can be obtained by setting $x' = 0$ and $y' = 0$. For the system (18) we obtain five fixed points (see Table 1). The coordinate of the point $\mathcal{A}$ is independent of ξ and n and the ones of $\mathcal{B}$ and $\mathcal{C}$ are independent of n . The point $\mathcal{D}$ is a finite fixed point for $n \neq 2$ and $\mathcal{E}$ a finite fixed point for $\xi \neq 1/2(n+2)$.

Merging occurs for the points $\mathcal{E}$ and $\mathcal{B}$, and $\mathcal{E}$ and $\mathcal{C}$ for $n = 4 \mp \frac{\sqrt{3}\sqrt{\xi(12\xi-1)}}{\xi}$ respectively. The point $\mathcal{E}$ also merges with $\mathcal{D}$ for $n = 4 \pm \sqrt{\frac{12\xi-1}{\xi}}$.

All the fixed points except $\mathcal{A}$ and $\mathcal{D}$ are associated with flat spatial geometry. Point $\mathcal{A}$ is associated with an open spatial geometry and for $\mathcal{D}$ the sign of the space curvature depends on ξ and n : $K_{\mathcal{D}}$ is positive for $n \neq 2, 4 - \sqrt{\frac{12\xi-1}{\xi}} < n < 4 + \sqrt{\frac{12\xi-1}{\xi}}$ and negative otherwise.

The stability of the fixed points can be determined by evaluating the eigenvalues of the Jacobian matrix associated with the system (18) (see Table 3), as prescribed by the Hartman-Grobman Theorem [39] §.

The fixed point $\mathcal{A}$ is a saddle for every value of the parameters ξ and n . The point $\mathcal{B}$ can either be a stable node or a saddle node whereas $\mathcal{C}$ is either an unstable node or a saddle node, depending on the values of ξ and n . The eigenvalues of $\mathcal{D}$ and $\mathcal{E}$ are both dependent on ξ and n so that the stability varies over the different ranges of ξ and n . The stability of these fixed points have been summarized in Table 4.

The coordinates of the fixed points may be used to determine exact cosmological solutions at the fixed points themselves. In fact, when evaluated at these points, the Friedmann and the Klein-Gordon equations can be written as:

$$\dot{H} = -\frac{H^2}{\alpha}, \quad \alpha = \left[1 - 2x_i - \frac{1}{6\xi}x_i^2 + \left(\frac{6\xi(n-2)-1}{12\xi-1} \right) y_i \right]^{-1}, \quad (19)$$

$$\frac{\ddot{\phi}}{\phi} = -\beta H^2, \quad \beta = 3x_i + x_i^2 + \frac{6\xi(n-4)}{1-12\xi}y_i, \quad (20)$$

where (x_i, y_i) represent the coordinates of the fixed points. Integrating (19) and substituting into (20) gives

$$\frac{\ddot{\phi}}{\phi} + \frac{\alpha^2\beta}{(t-t_0)^2} = 0, \quad (21)$$

which has a Cauchy-Euler form. If the terms α and β are different from zero, equations (19) and (21) can be easily integrated, giving

$$a = a_0 (t - t_0)^\alpha, \quad (22)$$

§ The values of the parameter for which the eigenvalues are zero are bifurcations for the dynamical system. In this paper we will not give an analysis of the bifurcations referring the reader to the specific literature for more details.

$$\phi = \begin{cases} (t - t_0)^{1/2} [\phi_0 (t - t_0)^m + \phi_1 (t - t_0)^{-m}] , & \text{if } \alpha^2 \beta < \frac{1}{4} , \\ (t - t_0)^{1/2} (\phi_0 + \phi_1 \ln t) , & \text{if } \alpha^2 \beta = \frac{1}{4} , \\ (t - t_0)^{1/2} (\phi_0 \sin [m \ln (t - t_0)] + \phi_1 \cos [m \ln (t - t_0)]) , & \text{if } \alpha^2 \beta > \frac{1}{4} , \end{cases} \quad (23)$$

where

$$m = \frac{1}{2} \sqrt{1 - 4\alpha^2 \beta}. \quad (24)$$

In the case of point $\mathcal{A}$, we have a Milne evolution and a constant scalar field

$$\alpha_{\mathcal{A}} = 1, \quad \phi_{\mathcal{A}} = \phi_0. \quad (25)$$

The above solution (as well as some others that will follow) is particularly interesting because the scalar field is *constant*. This implies that $\mathcal{A}$ represents a state in which G_{eff} is constant and the potential of the scalar field acts as a cosmological constant. In other words, at this point, scalar tensor gravity is indistinguishable from standard GR plus cosmological constant.

For the point $\mathcal{B}$ the solutions are given by (22) and (23) with

$$\alpha_{\mathcal{B}} = \frac{1}{3(1 - 8\xi) - 4\sqrt{3\xi(12\xi - 1)}}, \quad m_{\mathcal{B}} = \frac{3}{6(1 - 8\xi) - 8\sqrt{3\xi(12\xi - 1)}}, \quad (26)$$

and for point $\mathcal{C}$

$$\alpha_{\mathcal{C}} = \frac{1}{3(1 - 8\xi) + 4\sqrt{3\xi(12\xi - 1)}}, \quad m_{\mathcal{C}} = \frac{3}{6(1 - 8\xi) + 8\sqrt{3\xi(12\xi - 1)}}. \quad (27)$$

Note that, for $\xi < 0$, the value of α for the above two solutions is always positive and less than 1 i.e. these two solutions always represent two Friedmann-like solutions. In addition, $\mathcal{B}$ represents a solution in which the scalar field is growing, while at $\mathcal{C}$, ϕ is dissipating. If $n \neq 2$, the point $\mathcal{D}$ is associated with

$$\alpha_{\mathcal{D}} = 1, \quad m_{\mathcal{D}} = \frac{n + 2}{2(n - 2)}, \quad (28)$$

which represents another Milne solution, while the scalar field is decreasing for $n > 2$ and increasing for $n < 2$. It is interesting that, unlike the Milne solution in GR, this linear solution for a is not necessarily a spatially hyperbolic one. The constant a_0 can be related to the parameters ξ and n for non-flat solutions:

$$a_0^2 = -\frac{k(n - 2)^2 \xi}{1 + [4 + n(n - 8)]\xi}. \quad (29)$$

When $n = 2$, $\mathcal{D}$ becomes an asymptotic fixed point and merges with $\mathcal{D}_{\infty}$ (its solution will be presented within the asymptotic analysis). Finally, for point $\mathcal{E}$ we have

$$\alpha_{\mathcal{E}} = \frac{2(n + 2)\xi - 1}{(n - 4)(n - 2)\xi}, \quad m_{\mathcal{E}} = \frac{n + 2}{2(n - 2)} \quad (30)$$

for all $n \neq 2, 4$. This solution represents an expansion for $\frac{1 - 4\xi}{2\xi} < n < 2$ and $n > 4$ and was already found in other contexts [21–25]. In particular, in [25] it is shown that for $2 < n < 4$ the scale factor associated to this point evolves towards a superinflating state (also called “Big Rip” singularity) without including any exotic feature like ghosts or

Table 1. The coordinates and scale factor solutions of the fixed points for the vacuum case. We only show the exponent α of the solutions (22).

Point	Coordinates (x, y)	K	α
$\mathcal{A}$	$(0, 0)$	-1	1
$\mathcal{B}$	$(2(-6\xi - \sqrt{3\xi(12\xi - 1)}), 0)$	0	$\frac{1}{3(1-8\xi) - 4\sqrt{3\xi(12\xi - 1)}}$
$\mathcal{C}$	$(2(-6\xi + \sqrt{3\xi(12\xi - 1)}), 0)$	0	$\frac{1}{3(1-8\xi) + 4\sqrt{3\xi(12\xi - 1)}}$
$\mathcal{D}$	$(-\frac{2}{n-2}, \frac{2-24\xi}{3\xi(n-2)^2})$	$\frac{-1-[4+n(n-8)]\xi}{(n-2)^2\xi}$	$1, \left(a_0^2 = -\frac{k(n-2)^2\xi}{1+[4+n(n-8)]\xi}\right)$
$\mathcal{E}$	$(\frac{2(n-4)\xi}{1-2(n+2)\xi}, \frac{(12\xi-1)[3+(n-10)(n+2)\xi]}{3(1-2(n+2)\xi)^2})$	0	$\begin{cases} \frac{2(n+2)\xi-1}{(n-4)(n-2)\xi}, & n \neq 2, 4 \\ a = a_0, & n = 2 \\ a = a_0 e^{C(t-t_0)}, & n = 4 \end{cases}$

non standard fluids. When $n = 2$ and $n = 4$, using the cosmological equations we obtain the solutions

$$a = a_0, \quad \phi = \phi_0 \quad (\lambda = 0), \quad (31)$$

$$a = a_0 e^{C(t-t_0)}, \quad \phi = \phi_0 = \pm \sqrt{\frac{-6\xi C^2}{\lambda}}, \quad (32)$$

which represent a static universe and a de Sitter evolution respectively. In both these cases an effective cosmological constant is present, whose value depends on the effective gravitational constant (via ϕ_0) and the coupling constant λ of the self-interaction of the scalar field. Again, since the scalar field is constant in these cases, these solution are indistinguishable from the GR solutions. The difference with point $\mathcal{A}$ is that the solution does not occur in “pure” GR and can also be stable. This is particularly interesting in the second case ($n = 4$) in which a de-Sitter solution able to mimic an inflationary or dark energy phase in a Λ GR cosmology is a semi-global attractor. The possibility that scalar tensor gravity could converge to GR has been proposed within the context of extended inflation first using numerical techniques [40] and then with a more formal proof [41]. The dynamical system approach allows one to see this phenomenon in a very clear way, even in the more general case of a non-zero potential. It turns out that the nature of the potential, determined in our case by the value of n , together with the value of the coupling plays a critical role in the realization of this mechanism.

It is useful to define the deceleration parameter q in terms of the dynamical variables:

$$q = -\frac{\dot{H}}{H^2} - 1 = 2x_i + \frac{1}{6\xi}x_i^2 - \left(\frac{1+12\xi-6\xi n}{1-12\xi}\right)y_i. \quad (33)$$

This equation represents a parabola in the phase space that divides the accelerating ($q < 0$) expansion phases from the decelerating ($q > 0$) ones. Points $\mathcal{A}$ and $\mathcal{D}$ lie on the curve (33) as expected by the form of their scale factor solutions. On the other hand $\mathcal{B}$ and $\mathcal{C}$ always lie on the decelerated expansion side of the curve. Instead, for $\mathcal{E}$ we have decelerated expansion for $4 - \sqrt{\frac{12\xi-1}{\xi}} < n < 2$ or $4 < n < 4 + \sqrt{\frac{12\xi-1}{\xi}}$ and a decelerated one for $\frac{1-4\xi}{2\xi} < n < 4 - \sqrt{\frac{12\xi-1}{\xi}}$ or $n > 4 + \sqrt{\frac{12\xi-1}{\xi}}$.

Table 2. Values of the parameter m and the corresponding scalar field solutions for the fixed points in vacuum.

Point	m	Solutions
$\mathcal{A}$	$\frac{1}{2}$	$\phi = \phi_1, \quad \phi_0, \lambda = 0$
$\mathcal{B}$	$\frac{3}{6(1-8\xi)-8\sqrt{3\xi(12\xi-1)}}$	$\phi = \phi_0(t-t_0)^{1/2+m}, \quad \phi_1, \lambda = 0$
$\mathcal{C}$	$\frac{3}{6(1-8\xi)+8\sqrt{3\xi(12\xi-1)}}$	$\phi = \phi_1(t-t_0)^{1/2-m}, \quad \phi_0, \lambda = 0$
$\mathcal{D}$	$\frac{n+2}{2(n-2)}$	$\begin{cases} \phi = \phi_0^{\mathcal{D}}(t-t_0)^{1/2}, & n = -2 \\ \phi = \phi_1^{\mathcal{D}}(t-t_0)^{1/2-m}, & n \neq -2, 2 \\ \phi_0^{\mathcal{D}} = \pm \left(\frac{4\lambda}{1-12\xi} \right)^{\frac{1}{4}}, & \phi_1^{\mathcal{D}} = \left(\frac{4(1-12\xi)}{\lambda(n-2)^2} \right)^{\frac{1}{n-2}} \end{cases}$
$\mathcal{E}$	$\frac{n+2}{2(n-2)}$	$\begin{cases} \phi = \phi_0^{\mathcal{E}}(t-t_0)^{1/2}, & n = -2 \\ \phi = \phi_1^{\mathcal{E}}(t-t_0)^{1/2-m}, & n \neq -2, 2, 4 \\ \phi_0^{\mathcal{E}} = \pm 2 \left(\frac{6\lambda\xi}{12\xi-1} \right)^{\frac{1}{4}}, & \\ \phi_1^{\mathcal{E}} = \left(\frac{2(12\xi-1)[3+(n-10)n+2]\xi}{\lambda\xi(n-2)(n-4)} \right)^{\frac{1}{n-2}} \end{cases}$

Table 3. The eigenvalues associated with the fixed points in the vacuum model.

Point	Eigenvalues
$\mathcal{A}$	$[-2, 2]$
$\mathcal{B}$	$\left[4(1-12\xi) - 8\sqrt{3\xi(12\xi-1)}, 6-2(n+2) \left(6\xi + \sqrt{3\xi(12\xi-1)} \right) \right]$
$\mathcal{C}$	$\left[4(1-12\xi) + 8\sqrt{3\xi(12\xi-1)}, 6-2(n+2) \left(6\xi - \sqrt{3\xi(12\xi-1)} \right) \right]$
$\mathcal{D}$	$\left[\frac{(4-n)\xi - \sqrt{\xi\{3(8-n)n\xi-4\}}}{(n-2)\xi}, \frac{(4-n)\xi + \sqrt{\xi\{3(8-n)n\xi-4\}}}{(n-2)\xi} \right]$
$\mathcal{E}$	$\left[\frac{3+(n-10)(n+2)\xi}{2(n+2)\xi-1}, \frac{2+2\{4+(n-8)n\}\xi}{2(n+2)\xi-1} \right]$

4.2. Asymptotic analysis

Since the dynamical system (18) is not compact, it might admit an asymptotic structure that is relevant for the global dynamics. In order to analyze the asymptotic features of the phase space, we use the Poincaré projection [21, 42]. This method consists of transforming to the polar coordinates

$$x = \bar{r} \cos \psi, \quad y = \bar{r} \sin \psi \quad (34)$$

and setting $\bar{r} = \frac{\sqrt{r}}{1-r}$. In this way, the asymptotic regime is achieved for $r \rightarrow 1$. Using the Poincaré projection, the asymptotic form of the dynamical equations (18) read

$$r' = \frac{\cos^2 \psi (\cos 2\psi - 3)}{12\xi}, \quad (35)$$

Table 4. Stability of the fixed points in the vacuum case. The parameters are $N_{\pm} = 4 \pm \sqrt{3(12\xi - 1)/\xi}$, $P_{\pm} = 4 \pm 2\sqrt{(12\xi - 1)/3\xi}$ and $Q_{\pm} = 4 \pm \sqrt{(12\xi - 1)/\xi}$. We use the term ‘attractor’ to denote a sink, ‘repeller’ to denote a source and ‘spiral’ to denote an attractive spiral.

Points	$n < \frac{1-4\xi}{2\xi}$	$\frac{1-4\xi}{2\xi} \leq n < N_-$	$N_- < n < P_-$	$P_- < n < Q_-$
$\mathcal{A}$	saddle	saddle	saddle	saddle
$\mathcal{B}$	repeller	repeller	repeller	repeller
$\mathcal{C}$	saddle	saddle	repeller	repeller
$\mathcal{D}$	spiral	spiral	spiral	attractor
$\mathcal{E}$	attractor	repeller	saddle	saddle
	$Q_- < n < 2$	$2 < n < Q_+$	$Q_+ < n \leq P_+$	$P_+ < n < N_+$
$\mathcal{A}$	saddle	saddle	saddle	saddle
$\mathcal{B}$	repeller	repeller	repeller	repeller
$\mathcal{C}$	repeller	repeller	repeller	repeller
$\mathcal{D}$	saddle	saddle	attractor	spiral
$\mathcal{E}$	attractor	attractor	saddle	saddle
	$n > N_+$			
$\mathcal{A}$	saddle			
$\mathcal{B}$	saddle			
$\mathcal{C}$	repeller			
$\mathcal{D}$	spiral			
$\mathcal{E}$	repeller			

$$\psi' = -\frac{\cos^3 \psi \sin \psi}{6\xi(1-r)^2}. \quad (36)$$

Note that the radial equation does not depend on the radial coordinate. This means that the fixed points can be obtained by only considering the equation for ψ' . Setting $\psi' = 0$ we obtain the four fixed points listed in Table 5.

Let us now derive the behaviour of the scale factor at these points. This can be done by integrating the Friedmann equation (19) in the above asymptotic limit. The points $\mathcal{A}_{\infty}$ and $\mathcal{C}_{\infty}$ correspond to $y \rightarrow 0$ as $x \rightarrow \pm\infty$ respectively. In this limit the x' equation (18) reduces to

$$x' = -\frac{1}{6\xi} x^3, \quad (37)$$

which admits the solution

$$x^2 = \frac{3\xi}{\tau - \tau_{\infty}}. \quad (38)$$

In the given limit, the evolution equation (19) can be written as

$$\frac{H'}{H} = \frac{1}{6\xi} x^2, \quad (39)$$

which may be solved together with (38) to yield the solution

$$|\tau - \tau_{\infty}| = [C_1 \pm C_2(t - t_0)]^2, \quad (40)$$

which represents a universe which reaches a maximum radius and then recollapses. The points $\mathcal{B}_\infty$ and $\mathcal{D}_\infty$, instead, correspond to $x \rightarrow 0$ as $y \rightarrow \pm\infty$. In this limit the second of (18) reduces to

$$y' = \frac{2(6\xi(n-2)-1)}{12\xi-1} y^2, \quad (41)$$

which admits the solution

$$y = \frac{12\xi-1}{2(1-6\xi(n-2))(\tau-\tau_\infty)}. \quad (42)$$

Equation (19) now reduces to

$$\frac{H'}{H} = \frac{(1-6\xi(n-2))}{12\xi-1} y, \quad (43)$$

and substituting the solution for y , we obtain

$$|\tau - \tau_\infty| = [C_1 \pm C_2(t - t_0)]^2. \quad (44)$$

The stability of the asymptotic fixed points is summarized in Table 5. The points $\mathcal{A}_\infty$ and $\mathcal{C}_\infty$ are saddles for every value of n and ξ . The points $\mathcal{B}_\infty$ and $\mathcal{D}_\infty$ are non-hyperbolic and they can be shown to represent saddle-nodes. This means that they behave like saddles or nodes depending on which direction the orbits approach them and that a local separatrix exists to divide the different stability domains. In our specific case this separatrix corresponds to the equator of the Poincaré sphere (i.e. our “infinity”), so that effectively $\mathcal{B}_\infty$ behaves as a saddle if $n < 2 + 1/6\xi$ or $n = 2$ and an attractor if $n > 2 + 1/6\xi$ ($n \neq 2$), and $\mathcal{D}_\infty$ behaves like a saddle if $n > 2 + 1/6\xi$ (including $n = 2$) and an attractor if $n < 2 + 1/6\xi$. For both these point the separatrix (i.e. our unitary circle) is always attractive so that orbits can bounce off the saddle and then approach the point along the unitary circle (see for example Figure 1). A summary of the stability of the asymptotic fixed points is summarized in Table 5.

The behaviour of the scalar field can be obtained in the same way by substituting the solutions for x and y in the appropriate limit of equation (20). For the points $\mathcal{A}_\infty$ and $\mathcal{C}_\infty$ (20) we obtain

$$\phi = c_1 \cos\left(c_0 \sqrt{3\xi}(t - t_0)\right) + c_2 \sin\left(c_0 \sqrt{3\xi}(t - t_0)\right), \quad (45)$$

where c_0, c_1 and c_2 are integration constants. For the points $\mathcal{B}_\infty$ and $\mathcal{D}_\infty$ we obtain

$$\phi = c_1 \cos\left(c_0(t - t_0) \sqrt{\left|\frac{6\xi(n-4)}{2(1-6\xi(n-4))}\right|}\right) + c_2 \sin\left(c_0(t - t_0) \sqrt{\left|\frac{6\xi(n-4)}{2(1-6\xi(n-4))}\right|}\right), \quad (46)$$

for $n > 2 + \frac{1}{6\xi}$ or $n > 4$, and

$$\phi = c_1 e^{c_0(t-t_0) \sqrt{\frac{6\xi(n-4)}{2(1-6\xi(n-4))}}} + c_2 e^{c_0(t-t_0) \sqrt{\frac{6\xi(n-4)}{2(1-6\xi(n-4))}}}, \quad (47)$$

for $2 + \frac{1}{6\xi} < n < 4$.

Since the phase space is two dimensional we can easily draw phase space diagrams for the vacuum case. Here we will limit ourselves to four examples representing the global phase space || for four specific values of (n, ξ) (see Figures 1-4) that includes the two cases in which the theory admits a GR attractor.

|| By “global phase space” we mean the projection of the $\theta > \pi/2$ part of the Poincaré sphere on the plane that contains its equator.




Figure 1. Global phase space for $\frac{1-4\xi}{2\xi} < n < 4 - \sqrt{3}\sqrt{\frac{12\xi-1}{\xi}}$ (e.g. $n = -3$ and $\xi = -0.35$). Note that the non hyperbolic point B_∞ correspond to a saddle everywhere but on the border of the circle in which it behaves as an attractor (see the text for details).




Figure 2. Global phase space for $n = 2$ with $\xi = -1$.




Figure 3. Global phase space for $2 < n < 4 + \sqrt{\frac{12\xi-1}{\xi}}$ (e.g. $n = 4$ and $\xi = -1$).




Figure 4. Global phase space for $4 + \sqrt{\frac{12\xi-1}{\xi}} < n < 4 + \sqrt{3} \sqrt{\frac{12\xi-1}{\xi}}$ (e.g. $n = 8$ and $\xi = -1$).

Table 5. Coordinates, behaviour of the scale factor and stability of the asymptotic fixed points in the vacuum model.

Point	ψ	Behaviour	Stability
$\mathcal{A}_\infty$	0	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	saddle
$\mathcal{B}_\infty$	$\frac{\pi}{2}$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	$\begin{cases} \text{saddle} & \{n < 2 + \frac{1}{6\xi}\} + \{n = 2\} \\ \text{attractor} & \{n > 2 + \frac{1}{6\xi}\} - \{n = 2\} \end{cases}$
$\mathcal{C}_\infty$	π	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	saddle
$\mathcal{D}_\infty$	$\frac{3\pi}{2}$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	$\begin{cases} \text{attractor} & n < 2 + \frac{1}{6\xi} \\ \text{saddle} & n > 2 + \frac{1}{6\xi} \end{cases}$

5. The matter case

As we have seen in the system (14), the presence of matter in the dynamical system equations implies the introduction of another variable: z . Since μ is defined to be positive, the definition of the dynamical variables tells us that only negative values of z are compatible with attractive gravity. For this reason we will restrict the following analysis only to the physically relevant case $z \leq 0$.

As in the vacuum case, we can use the constraint (15) to eliminate one of the dynamical variables. The system (14) then reduces to:

$$\begin{aligned}
x' &= \frac{1}{12\xi - 1} \left[2(1 - 12\xi)x + 4(1 - 12\xi)x^2 - \left(2 - \frac{1}{6\xi}\right)x^3 + 6\xi(n - 4)y \right. \\
&\quad \left. - (1 - 6\xi(n - 2))xy - \frac{1}{2}(1 + 3w - 24\xi)xz + 6\xi(1 - 3w)z \right], \\
y' &= \frac{y}{12\xi - 1} \left[2(12\xi - 1) + (12\xi - 1)(n - 6)x + 2\left(2 - \frac{1}{6\xi}\right)x^2 \right. \\
&\quad \left. - 2(1 - 6\xi(n - 2))y - (1 + 3w - 24\xi)z \right], \\
z' &= \frac{z}{12\xi - 1} \left[(1 + 3w)(12\xi - 1) - 6(12\xi - 1)x \right. \\
&\quad \left. + 2\left(2 - \frac{1}{6\xi}\right)x^2 - 2(1 - 6\xi(n - 2))y - (1 + 3w - 24\xi)z \right].
\end{aligned} \tag{48}$$

Since $y' = 0$ and $z' = 0$ are zero for $y = 0$ and $z = 0$, the two planes $y = 0$ and $z = 0$ corresponds to two invariant submanifolds. The first plane represents classes of theories in which the potential is zero, the second one classes of theories for which $z = 0$ and constitutes part of a vacuum invariant submanifold. The structure of the total vacuum invariant submanifold can be derived by writing the energy density in terms of the dynamical variables [31, 32]:

$$\mu \propto zy^{\frac{2}{n-2}} H^{\frac{2n}{n-2}}. \tag{49}$$

It is clear that when $z = 0$ and $y \neq 0$ the energy density is zero. However when $y = 0$ and $z \neq 0$ the behaviour of μ depends on the value of n : the energy density is zero when $n > 2$, but it is divergent when $n < 2$. When both y and z are equal to zero, we can conclude that $\mu = 0$ only if $n > 2$. For $n < 2$ (49) is divergent and μ can only be obtained by directly solving the field equations.

5.1. Finite analysis

Setting $x' = 0$, $y' = 0$ and $z' = 0$ we obtain eight fixed points (see Table 6). The first five ($\mathcal{A}$, $\mathcal{B}$, $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$) sit in the $z = 0$ plane and have the same (x, y) coordinates of the corresponding vacuum fixed points. The coordinates of $\mathcal{F}$ and $\mathcal{G}$ are independent of n and are finite for all ξ ; $\mathcal{H}$ is a finite fixed point for $n \neq 0$ and merges with the asymptotic fixed subspace $\mathcal{L}_\infty$ for $n = 0$.

Merging occurs between the points $\mathcal{B}$, $\mathcal{C}$, $\mathcal{D}$, $\mathcal{E}$, $\mathcal{G}$ and $\mathcal{H}$. The first four points merge in the same way and for the same values of the parameters given in the vacuum case. Point $\mathcal{H}$ merges with $\mathcal{E}$ for $n = \frac{1}{2} \left(7 + 3w + \sqrt{\frac{\xi(9w^2 + 66w + 73) - 6(w+1)}{\xi}} \right)$; and with $\mathcal{G}$ when $n = \frac{3(w+1)(8\xi+w-1)}{4\xi(3w-1)}$.

All the fixed points but $\mathcal{A}$, $\mathcal{D}$, $\mathcal{F}$ are associated with flat solutions. $\mathcal{A}$ is associated with negative curvature and the sign of the spatial curvature for point $\mathcal{D}$ and $\mathcal{F}$ depends on the value of n and ξ .

The stability of the fixed points may be determined using the Hartman-Grobman theorem as in the vacuum case. The point $\mathcal{A}$ is a saddle for every value of w . Point $\mathcal{B}$ keeps the same stability found in the vacuum case for $w = 0$ and $1/3$, but it is always a saddle when $w = 1$. Instead, the stability of points $\mathcal{C}$ and $\mathcal{D}$ is the same as in the vacuum case for every value of w . Point $\mathcal{E}$ varies its stability with ξ, n, w and can behave like an attractor, a repeller or a saddle as shown in Table 10. On the other hand, the fixed points $\mathcal{F}$ and $\mathcal{G}$ are saddles for every value of the parameters. Finally, $\mathcal{H}$ is a saddle or a saddle focus for $w = 0$ and $w = 1/3$, while for $w = 1$ it can also be a repeller or an anti-spiral. Its stability is summarized in Table 11.

As in the vacuum case the coordinates of the fixed points can be used to find exact solutions for the evolution of the scale factor. From (9) and (11) we have

$$\dot{H} + \frac{H^2}{\alpha_m} = 0, \quad \alpha_m = \left[1 - 2x_i - \frac{1}{6\xi}x_i^2 + \left(\frac{1+12\xi-6\xi n}{1-12\xi} \right) y_i + \left(\frac{1+3w-24\xi}{2-24\xi} \right) z_i \right]^{-1} \quad (50)$$

$$\frac{\ddot{\phi}}{\phi} + \frac{\alpha_m^2 \beta}{(t-t_0)^2} = 0, \quad \beta = 3x_i + x_i^2 + \frac{6\xi(n-4)}{1-12\xi}y_i + \frac{6\xi(1-3w)}{1-12\xi}z_i, \quad (51)$$

where (x_i, y_i, z_i) represents the coordinates of the fixed points. For α and β different from zero, these equations have solutions of the form (22) and (23).

The points $\mathcal{A}$, $\mathcal{B}$, $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ have the same solutions as in the vacuum case, since for these points $z = 0$. In particular the convergence mechanism of scalar tensor gravity towards GR is preserved when matter is present. Such a result was expected since in [40] and [41] this phenomenon is described when matter is present. The point $\mathcal{F}$ is associated to the Milne evolution and a constant scalar field

$$\alpha_{\mathcal{F}} = 1 \quad \phi_{\mathcal{F}} = \phi_0, \quad (52)$$

the value of the constant a_0 being influenced by the parameters ξ and w for non-zero spatial curvature

$$a_0^2 = \frac{16k\xi}{1 - 16\xi + (2 - 3w)w}. \quad (53)$$

However, a direct check of the equations reveals that this solution is valid only for $w = -1/3$. Since in our model standard matter For point $\mathcal{G}$ we have

$$\alpha_{\mathcal{G}} = \frac{2(8\xi - 1 + w)}{32\xi + 3(w^2 - 1)}, \quad m_{\mathcal{G}} = \frac{3w^2 + 48\xi w + 16\xi - 3}{2(3w^2 + 32\xi - 3)}. \quad (54)$$

This solution represents a Friedmann like expansion with the exponent depending on both ξ and w . For $0 < w < 1/3$ we have $\frac{1}{2} \leq \alpha_G < \frac{2-16\xi}{3-32\xi}$, for $1/3 \leq w < 1$ we have $\frac{1}{3(1-8\xi) - 4\sqrt{3\xi(12\xi-1)}} \leq \alpha_G < \frac{1}{2}$. Direct comparison with the cosmological equations reveals that this solution is physical only for $w > 2/3$ and $w \neq 1$.

Finally, point $\mathcal{H}$ with $n \neq 2$ is linked to a solution that a first glance resembles a well known Friedmann GR one:

$$\alpha_{\mathcal{H}} = \frac{2n}{3(n-2)(1+w)}, \quad m_{\mathcal{H}} = \frac{n+2}{2(n-2)}. \quad (55)$$

This solution represents a Friedmann-like expansion for $n < 0$ and $n > \frac{6(w+1)}{3w+1}$, a contraction for $0 < n < 2$ and power-law inflation when $2 < n < \frac{6(w+1)}{3w+1}$. When $n = 2$ we obtain a solution of the form

$$a = a_0, \quad \phi = \phi_0, \quad \lambda = 0, \quad (56)$$

which corresponds to a static universe with cosmological constant. Also this solution represent an effective “GR state” for the scalar tensor cosmology but in this case it is not a stable one. This means that in our model of non-vacuum scalar tensor cosmology no stable non vacuum GR like solutions are allowed.

Using (49) and the cosmological equations it is easy to conclude that the points $\mathcal{F}$ and $\mathcal{G}$ both admit vacuum solutions. This is not true for $\mathcal{H}$, for which the energy density is

$$\mu = \mu_0 t^{-\frac{2n}{n-2}}, \quad (57)$$

when $n \neq 2$, where

$$\mu_0 = z_{\mathcal{H}} \left(\frac{2y_{\mathcal{H}}}{\lambda} \right)^{\frac{2}{n-2}} \left[\frac{4\xi n^2}{3(n-2)^2(1+w)^2} \right]^{\frac{n}{n-2}},$$

and $y_{\mathcal{H}}$ and $z_{\mathcal{H}}$ are the y and z coordinates of $\mathcal{H}$ respectively. It is clear from the expression above that this point does not represent a physical solution for all values of n and ξ since μ_0 and therefore μ can, in principle, be negative. In order for us to determine the values of n and ξ for which $\mathcal{H}$ is physical, we have to solve the inequality $\mu_0 > 0$. We obtain

$$\frac{1}{2}(7+3w) - \sqrt{\frac{\xi(9w^2+66w+73)-6(w+1)}{4\xi}} < n < \frac{1}{2}(7+3w) + \sqrt{\frac{\xi(9w^2+66w+73)-6(w+1)}{4\xi}}. \quad (58)$$

When $n = 2$ the cosmological equations reveal that $\mathcal{H}$ is instead associated with a vacuum ($\mu = 0$) solution.

As in the vacuum case we can write the deceleration parameter q in terms of the dynamical variables via

$$q = -\frac{\dot{H}}{H^2} - 1 = 2x_i + \frac{1}{6\xi}x_i^2 - \left(\frac{1+12\xi-6\xi n}{1-12\xi} \right) y_i - \left(\frac{1+3w-24\xi}{2-24\xi} \right) z_i. \quad (59)$$

The surface defined above divides the phase space in two volumes representing the region in which the expansion is accelerated and the region in which it is decelerated. The fixed points $\mathcal{A}$, $\mathcal{B}$, $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ behave like their vacuum counterparts, since they all satisfy $z = 0$. Point $\mathcal{F}$ lies on this surface consistently with the fact that it corresponds to Milne evolution. The point $\mathcal{G}$ always lies in the region in which the

Table 6. The coordinates and the sign of the spatial curvature of the non vacuum fixed points.

Point	Coordinates (x, y, z)	K
$\mathcal{A}$	$(0, 0, 0)$	-1
$\mathcal{B}$	$\left(2(-6\xi - \sqrt{3\xi(12\xi - 1)}), 0, 0\right)$	0
$\mathcal{C}$	$\left(2(-6\xi + \sqrt{3\xi(12\xi - 1)}), 0, 0\right)$	0
$\mathcal{D}$	$\left(-\frac{2}{n-2}, \frac{2-24\xi}{3\xi(n-2)^2}, 0\right)$	$-\frac{1+[4+n(n-8)]\xi}{(n-2)^2\xi}$
$\mathcal{E}$	$\left(\frac{2(n-4)\xi}{1-2(n+2)\xi}, \frac{(12\xi-1)[3+(n-10)(n+2)\xi]}{3(1-2(n+2)\xi)^2}, 0\right)$	0
$\mathcal{F}$	$\left(-\frac{(1+3w)}{2}, 0, \frac{(12\xi-1)(1+3w)}{12\xi}\right)$	$\frac{1-16\xi+(2-3w)w}{16\xi}$
$\mathcal{G}$	$\left(\frac{4\xi(1-3w)}{8\xi-1+w}, 0, \frac{(12\xi-1)[3(w-1)^2-16\xi(2-3w)]}{3(8\xi-1+w)^2}\right)$	0
$\mathcal{H}$	$\left(-\frac{3(1+w)}{n}, \frac{3-3w^2-4\xi[6(1+w)+n(1-3w)]}{4n^2\xi}, \frac{3(1+w)+2\xi[n^2-6(1+w)-n(7+3w)]}{2n^2\xi}\right)$	0

Table 7. The exponent α of the scale factor solutions and the energy density for the non vacuum case.

Point	α	Matter density
$\mathcal{A}$	1	$\mu = 0$
$\mathcal{B}$	$\frac{1}{3(1-8\xi)-4\sqrt{3\xi(12\xi-1)}}$	$\mu = 0$
$\mathcal{C}$	$\frac{1}{3(1-8\xi)+4\sqrt{3\xi(12\xi-1)}}$	$\mu = 0$
$\mathcal{D}$	$1, \left(a_0^2 = -\frac{k(n-2)^2\xi}{1+[4+n(n-8)]\xi}\right)$	$\mu = 0$
$\mathcal{E}$	$\begin{cases} \frac{2(n+2)\xi-1}{(n-4)(n-2)\xi}, & n \neq 2, 4 \\ a = a_0 e^{C_0(t-t_0)}, & n = 2, 4 \end{cases}$	$\mu = 0$
$\mathcal{F}$	$1, \left(a_0^2 = \frac{16k\xi}{1-16\xi+(2-3w)w}\right)$	$\mu = 0$
$\mathcal{G}$	$\frac{2(8\xi-1+w)}{32\xi+3(w^2-1)}$	$\mu = 0$
$\mathcal{H}$	$\begin{cases} \frac{2n}{3(n-2)(1+w)}, & n \neq 2 \\ a = a_0, & n = 2 \end{cases}$	$\begin{cases} \mu = \mu_0 t^{-\frac{2n}{n-2}} \\ \mu = 0 \end{cases}$

Table 8. The parameter m and the corresponding scalar field solutions for the non vacuum case. The integration constants have been calculated by direct substitution in the cosmological equations.

Point	m	Solutions
$\mathcal{A}$	$\frac{1}{2}$	$\phi = \phi_1, \quad \lambda = 0$
$\mathcal{B}$	$\frac{3}{2(24\xi+4\sqrt{3\xi(12\xi-1)}-3)}$	$\phi = \phi_0(t-t_0)^{1/2+m}, \quad \lambda = 0$
$\mathcal{C}$	$\frac{3}{2(-24\xi+4\sqrt{3\xi(12\xi-1)}+3)}$	$\phi = \phi_1(t-t_0)^{1/2-m}, \quad \lambda = 0$
$\mathcal{D}$	$\frac{n+2}{2(n-2)}$	$\begin{cases} \phi = \phi_0^{\mathcal{D}}(t-t_0)^{1/2}, & n = -2 \\ \phi = \phi_1^{\mathcal{D}}(t-t_0)^{1/2-m}, & n \neq -2, 2 \end{cases}$
$\mathcal{E}$	$\frac{n+2}{2(n-2)}$	$\begin{cases} \phi = \phi_0^{\mathcal{E}}(t-t_0)^{1/2}, & n = -2 \\ \phi = \phi_1^{\mathcal{E}}(t-t_0)^{1/2-m}, & n \neq -2, 2 \end{cases}$
$\mathcal{F}$	$\frac{1}{2}(3w+2)$	$\phi = \phi_1 \ (w = -1/3), \quad \lambda = 0$
$\mathcal{G}$	$\frac{3w^2+48\xi w+16\xi-3}{2(3w^2+32\xi-3)}$	$\lambda = 0, \quad \xi = \frac{3(w-1)^2}{16(3w-2)} \rightarrow (w > 2/3, w \neq 1)$
$\mathcal{H}$	$\frac{n+2}{2(n-2)}$	$\begin{cases} \phi = \phi_0^{\mathcal{H}}(t-t_0)^{1/2}, & n = -2 \\ \phi = \phi_1^{\mathcal{H}}(t-t_0)^{1/2-m}, & n \neq -2, 2 \\ \phi_0^{\mathcal{H}} = \pm 2 \left(\frac{24\lambda(1+w)^2}{3(1-w^2)-16\xi(1+3w)} \right)^{\frac{1}{4}}, \\ \phi_1^{\mathcal{H}} = \left(\frac{2(3-4(n+6)+12(n-2)\xi w-3w^2)}{3\lambda(n-2)^2(1+w)^2} \right)^{\frac{1}{n-2}} \end{cases}$

expansion is decelerating. The behaviour of $\mathcal{H}$ depends on the barotropic factor w and the parameter n : it lies in the accelerated evolution region for $w = 0$ when $n < 0$ and $n > 6$, for $w = 1/3$ when $n < 0$ and $n > 4$ and for $w = 1$ when $n < 0$ and $n > 3$. Otherwise it lies in the decelerated evolution region.

5.2. Asymptotic analysis

We next study the asymptotic behaviour of the system (48) using the Poincaré projection. The compactification of the phase space can be achieved by transforming to spherical coordinates

$$x = \bar{r} \sin \theta \cos \psi, \quad y = \bar{r} \sin \theta \sin \psi, \quad z = \bar{r} \cos \theta, \quad (60)$$

where $\bar{r} = \frac{\sqrt{r}}{1-r}$ and $\bar{r} \in [0, \infty)$, $\theta \in [0, \pi]$ and $\psi \in [0, 2\pi]$. In the limit $r \rightarrow 1$ ($\bar{r} \rightarrow \infty$), equations (48) become

$$r' = \frac{\cos^2 \psi \sin^2 \theta}{24\xi} [(\cos 2\psi - 3) \sin^2 \theta - 4 \cos^2 \theta], \quad (61)$$

$$\theta' = -\frac{\cos \theta \sin^3 \theta \cos^4 \psi}{6\xi(1-r)^2}, \quad (62)$$

$$\psi' = \frac{\sin^2 \theta \cos^3 \psi \sin \psi}{6\xi(1-r)^2}. \quad (63)$$

Table 9. The eigenvalues associated with the non vacuum fixed points.

Point	Eigenvalues
$\mathcal{A}$	$[-2, 2, -1 - 3w]$
$\mathcal{B}$	$\left[4(1 - 12\xi) - 8\sqrt{3\xi(12\xi - 1)}, 6 - 2(n + 2) \left(6\xi + \sqrt{3\xi(12\xi - 1)} \right), \right.$ $\left. 3(1 - 8\xi) - \sqrt{3\xi(12\xi - 1)} - 3w \right]$
$\mathcal{C}$	$\left[4(1 - 12\xi) + 8\sqrt{3\xi(12\xi - 1)}, 6 - 2(n + 2) \left(6\xi - \sqrt{3\xi(12\xi - 1)} \right), \right.$ $\left. 3(1 - 8\xi) + \sqrt{3\xi(12\xi - 1)} - 3w \right]$
$\mathcal{D}$	$\left[\frac{(4-n)\xi - \sqrt{\xi\{3(8-n)n\xi - 4\}}}{(n-2)\xi}, \frac{(4-n)\xi + \sqrt{\xi\{3(8-n)n\xi - 4\}}}{(n-2)\xi}, \frac{6-n}{n-2} - 3w \right]$
$\mathcal{E}$	$\left[\frac{3+(n-10)(n+2)\xi}{2(n+2)\xi-1}, \frac{2+2\{4+(n-8)n\}\xi}{2(n+2)\xi-1}, \frac{3+2\{-6+(n-7)n\}\xi}{2(n+2)\xi-1} - 3w \right]$
$\mathcal{F}$	$\left[\frac{1}{2} (4 + (2 - n)(1 + 3w)), \frac{-2\xi(1-3w) - \sqrt{2\xi\{18\xi(1+w)^2 - (1-w)(1+3w)^2\}}}{4\xi}, \right.$ $\left. \frac{-2\xi(1-3w) + \sqrt{2\xi\{18\xi(1+w)^2 - (1-w)(1+3w)^2\}}}{4\xi} \right]$
$\mathcal{G}$	$\left[\frac{16\xi - (1-w)(1+3w)}{8\xi - 1 + w}, \frac{3(1-w)^2 - 16\xi(2-3w)}{2(8\xi - 1 + w)}, \frac{3(w^2 - 1) + 2\xi\{6+n+3(2-n)w\}}{8\xi - 1 + w} \right]$
$\mathcal{H}$	$\left[\frac{-4+(n+2)(1+3w)}{n}, \frac{3(w-1)(n+2)+12-\sqrt{S(n,\xi,w)}}{4n}, \frac{3(w-1)(n+2)+12+\sqrt{S(n,\xi,w)}}{4n} \right]$
$S(n, \xi, w) = 3\xi^{-1}(-1 + 12\xi)^{-1} \left\{ -36(w-1)(1+w)^2 + 3\xi [4(1+w)^2(12w-37) \right.$ $+ n^2(7 + (2-9w)w) + 4n(1+w)(w(19+6w)-17)] + 4\xi^2 [324(1+w)^2$ $+ 8n^3(-1+3w) + 12n(1+w)(29+3w) + n^2(17-21w(10+3w))] \}$	

Table 10. Stability of the fixed point $\mathcal{E}$ for the matter case. The parameters are $N_{\pm} = 4 \pm \sqrt{3}\sqrt{(12\xi-1)/\xi}$, $P_{\pm}^w = \frac{1}{2}(7+3w) \pm \sqrt{\frac{\xi(9w^2+66w+73)-6(1+w)}{4\xi}}$ and $Q_{\pm} = 4 \pm \sqrt{(12\xi-1)/\xi}$.

	$n < \frac{1-4\xi}{2\xi}$	$\frac{1-4\xi}{2\xi} < n < N_-$	$N_- < n < P_-$	$P_- < n < Q_-$
$w = 0, 1/3$	attractor	repeller	saddle	saddle
$w = 1$	attractor	repeller	saddle	saddle
	$Q_- < n < Q_+$	$Q_+ < n < P_+$	$P_+ < n < N_+$	$N_+ < n < P_+^1$
$w = 0, 1/3$	attractor	saddle	saddle	repeller
$w = 1$	attractor	saddle	saddle	saddle
	$n > P_+^1$			
$w = 0, 1/3$	repeller			
$w = 1$	repeller			

Table 11. Stability of the fixed point $\mathcal{H}$ of the matter case. The parameters are $P_{\pm} = \frac{1}{2}(7 + 3w) \pm \sqrt{\frac{\xi(9w^2 + 66w + 73) - 6(1+w)}{4\xi}}$ and S_i , which are the real roots of the polynomial $S(n, \xi, w)$ given in Table 9, for the given value of w . We use the term ‘anti-spiral’ to denote a pure repulsive spiral and ‘saddle focus’ to denote an unstable point with two complex eigenvalues.

$w = 0$	$n \leq S_1$	$S_1 < n < S_2$	$S_2 \leq n \leq S_3$	$n > S_3$
	saddle ($n \neq \frac{3(1-8\xi)}{4\xi}$)	saddle focus	saddle ($n \neq 0, 6, P_{\pm}$)	saddle focus
$w = 1/3$	$n \leq S_1$	$S_1 < n < S_2$	$n > S_2$	
	saddle focus	saddle ($n \neq 0, 4, P_{\pm}$)	saddle focus	
$w = 1$	$n \leq S_1$	$S_1 \leq n < P_+$	$P_+ < n \leq S_2$	$S_2 < n < S_3$
	saddle focus	saddle ($n \neq 0, 3, P_-$)	repeller	anti-spiral
	$S_3 \leq n < 6$	$n > 6$		
	repeller	saddle		

As in the vacuum case the radial equation does not contain the radial coordinate, so that the fixed points can be obtained using only the angular equations. Setting $\psi' = 0$ and $\theta' = 0$, we obtain four fixed points and a fixed subspace which are listed in Table 12.

The points $\mathcal{A}_{\infty}$ and $\mathcal{B}_{\infty}$ lie on the plane $z = 0$ and therefore their solutions are the same as the vacuum points $\mathcal{A}_{\infty}$ and $\mathcal{C}_{\infty}$, respectively. Points $\mathcal{C}_{\infty}$ and $\mathcal{D}_{\infty}$ represent the poles of the Poincaré sphere and it is easy to prove that they are linked to the same solutions at $\mathcal{A}_{\infty}$ and $\mathcal{B}_{\infty}$. The solution for the scalar field instead reads

$$\phi = c_1 e^{c_0(t-t_0)\sqrt{\frac{1+3w-24\xi}{2-24\xi}}} + c_2 e^{c_0(t-t_0)\sqrt{\frac{1+3w-24\xi}{2-24\xi}}}. \quad (64)$$

In addition to the fixed points above, we found two fixed subspaces $\mathcal{L}_1$ and $\mathcal{L}_2$, that contains all the points with $\psi = \pi/2$ and $\psi = 3\pi/2$, respectively. These subspaces corresponds to $x \rightarrow 0$, $y/z \rightarrow \pm \tan \theta_0$, respectively. The equations for y' and z' in this limit reduce to

$$y' = \frac{y^2}{1-12\xi} [2(1-6\xi(n-2)) \pm (1+3w-24\xi) \cot \theta_0], \quad (65)$$

$$z' = \frac{z^2}{1-12\xi} [(1+3w-24\xi) \pm 2(1-6\xi(n-2)) \tan \theta_0], \quad (66)$$

which admit the solutions

$$y = \frac{12\xi - 1}{(\tau - \tau_{\infty}) [2(1-6\xi(n-2)) \pm (1+3w-24\xi) \cot \theta_0]}, \quad (67)$$

$$z = \frac{12\xi - 1}{(\tau - \tau_{\infty}) [(1+3w-24\xi) \pm 2(1-6\xi(n-2)) \tan \theta_0]}. \quad (68)$$

In the limit above, (50) becomes

$$\dot{H} = - \left[\left(\frac{1+12\xi-6\xi n}{1-12\xi} \right) y + \left(\frac{1+3w-24\xi}{2-24\xi} \right) z \right] H^2, \quad (69)$$

which implies

$$|\tau - \tau_\infty| = [C_1 \pm C_2(t - t_0)]^2. \quad (70)$$

The solution for the scalar field in the fixed subspaces $\mathcal{L}_1$ and $\mathcal{L}_2$, can be found in similar way. In the limit above, using (67), (68) and (70) in (51) we find

$$\frac{\ddot{\phi}}{\phi} + A c_0^2 = 0, \quad (71)$$

where

$$A = \frac{6\xi(1-3w)\cos\theta_0 \pm 12\xi(n-4)\sin\theta_0}{2(1+3w-24\xi)\cos\theta_0 \pm 4(1-6\xi(n-2))\sin\theta_0}. \quad (72)$$

We find the following solutions

$$\phi = \begin{cases} c_1 \cos(\sqrt{|A|}c_0(t-t_0)) + c_2 \sin(\sqrt{|A|}c_0(t-t_0)), & \text{if } A < 0, \\ c_1 e^{\sqrt{A}c_0(t-t_0)} + c_2 e^{-\sqrt{A}c_0(t-t_0)}, & \text{if } A > 0. \end{cases} \quad (73)$$

The stability of these asymptotic fixed points can be deduced by analyzing the stability with respect to the angular coordinates, and from the sign of r' . The points $\mathcal{A}_\infty$ and $\mathcal{B}_\infty$ are stable nodes for all values of n and ξ . Points $\mathcal{C}_\infty$ and $\mathcal{D}_\infty$ instead are always saddles.

All the other points including the fixed subspaces $\mathcal{L}_1$ and $\mathcal{L}_2$, have both eigenvalues equal to zero. In order to derive the stability of these points we have to analyze the effect of the non-linear contributions have on dynamical equations. This can be done by Taylor developing the R.H.S of the dynamical equations around the fixed point up to the first non-zero order and then directly the resulting system.

For the points $\mathcal{C}_\infty$ and $\mathcal{D}_\infty$ we obtain the solutions:

$$\psi_{C_\infty} = c_2, \quad \theta_{C_\infty} = -\frac{6\xi}{\tau \sin(\psi_0) \cos^3(\psi_0) + 6\xi c_1}, \quad (74)$$

$$\psi_{D_\infty} = c_2, \quad \theta_{D_\infty} = \pi - \frac{6\xi}{\tau \sin(\psi_0) \cos^3(\psi_0) + 6\xi c_1}. \quad (75)$$

This result tells us that when we choose initial conditions around these points, the evolution of the universe will follow an orbit with constant ψ , with τ increasing $\P$ and θ approaching 0 or π . If we also consider the behaviour of the radial equation we conclude that these points behave like saddles.

For the fixed subspaces $\mathcal{L}_1$ and $\mathcal{L}_2$, we find

$$\psi(\tau) = \frac{\sqrt[3]{-2\xi}}{\sqrt[3]{\tau \cos\theta_0 \sin^3\theta_0 + 96\xi c_1}} \pm \frac{\pi}{2}, \quad (76)$$

$$\theta(\tau) = \theta_0 - \frac{\tan\theta_0}{2} + c_2 [\tau (\sin 4\theta_0 - 2 \sin 2\theta_0) - 768\xi c_1]^{\frac{2}{3} \csc^2 \theta_0}. \quad (77)$$

It is clear that, for τ increasing, the solution for ψ approaches $\pi/2$ for $\mathcal{L}_1$ and $3\pi/2$ for $\mathcal{L}_2$, while θ increases. The radial behaviour is very complicated and depends critically on the value of the coordinate θ_0 , w , n and ξ (we will not show this dependence here). We may however conclude that this subspace is a saddle or an attractor depending on the value of θ_0 .

$\P$ This behaviour is reversed in case of a contracting cosmology because in this case τ effectively changes sign when a is decreasing.

Table 12. Coordinates, behaviours and stability of the asymptotic fixed points in the non vacuum case. The value of the parameter A is given in (72).

Point	(θ, ψ)	Scale factor	Stability
$\mathcal{A}_\infty$	$(\frac{\pi}{2}, 0)$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	attractor
$\mathcal{B}_\infty$	$(\frac{\pi}{2}, \pi)$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	attractor
$\mathcal{C}_\infty$	$(0, 0)$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	saddle
$\mathcal{D}_\infty$	(π, π)	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	saddle
$\mathcal{L}_1$	$(\theta_0, \frac{\pi}{2})$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	
$\mathcal{L}_2$	$(\theta_0, \frac{3\pi}{2})$	$ \tau - \tau_\infty = [C_1 \pm C_2(t - t_0)]^2$	

The phase space in the matter case is 3-dimensional and therefore cannot be visualized as easily as its vacuum counterpart. For this reasons we will not give here any sketch of the phase space and we refer the reader to the next section for an analysis of the results derived above.

6. Discussion and conclusions

In this paper, the dynamics of STG FLRW cosmological models has been studied using a phase space analysis. We have considered a generic non-minimally coupled theory of gravity where the coupling and the potential are powers of the scalar field, both in a vacuum and in presence of a perfect fluid. The set of parameters characterizing the cosmological models are $\{\xi, \lambda, n, w\}$, i.e. the coupling constant, a constant parameter, the power of the self-interacting potential and the barotropic index of the perfect matter fluid, respectively. The phase-space is 3-dimensional in absence of matter while it is 4-dimensional in presence of matter, but in both cases the FLRW Hamiltonian constraint allows one to reduce their dimensionality. Our investigation considered the existence and local stability of critical points (finite analysis) and the asymptotic analysis via the Poincaré projection.

We identified five finite fixed points in the vacuum case and eight in the matter case. In the vacuum case, there are four asymptotic stability points corresponding to the four intersections of the axes with the unitary Poincaré circle. In the matter case, we have to consider a unitary 3-sphere and in additin to the vacuum asymptotic points we find two more points and two “fixed subspaces”. The stability of fixed points strictly depends on the values of the above parameters and in particular on ξ and n , i.e. the coupling and the power of self-interacting potential.

In the vacuum case, the two dimensional phase space is divided in two halves by the invariant submanifold associated with $V(\phi) = 0$. Of the nine fixed points we found, five of them ($\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{A}_\infty, \mathcal{C}_\infty$) are permanently on this invariant submanifold, two of them $\mathcal{B}_\infty, \mathcal{C}_\infty$ do not change their position and the other two “move” in the phase plane depending on the values of the parameters. In particular, $\mathcal{D}$ is characterized by $q = 0$ and its position is on the curve described by the equation given in (33) and $\mathcal{E}$ is associated with a flat spatial geometry and always lies on the curve described by the constraint (17) with $K = 0$. Even if matter is not present the scalar field is able to induce an expanding Friedmann-like evolution. However, the value of α in (26) and (27) reveals that at these fixed points the scale factor cannot grow faster

than $t^{1/2}$. In addition, these solutions are unstable for every value of the parameters. The point $\mathcal{E}$ admits the widest spectrum of behaviours. Depending on the values of n and ξ its solution can represent an inflationary phase, a Friedmann-like phase or a contraction. A comparison with the stability analysis reveals that $\mathcal{E}$ is an attractor only when it corresponds to contracting solutions or power law inflation. This has two consequences: (i) in the scalar field dominated regime our model of scalar tensor cosmology admits an inflationary phase as an attractor even if this attractor is not global and not unique; (ii) since there is no value of the parameters for which the $\mathcal{E}$ is a saddle, this model does not admit a transient inflationary phase and cannot solve the graceful exit problem.

The case $n = 4$ is particularly interesting because, for this value of n , $\mathcal{E}$ is associated with a GR ($\phi = \text{const.}$) de Sitter solution and is an attractor. This fact has two main consequences: (i) it shows that with a non-minimally coupled scalar field with quartic potential, the early time cosmology evolves towards an inflationary phase which is indistinguishable from a GR one; and (ii) it gives us an independent confirmation of the idea that a scalar tensor theory of gravity can evolve towards GR. The difference is that when a potential is present, the realization of such mechanism is strictly related to the form of the potential.

In the matter case, we found three new finite fixed points which are physical only for specific values of the parameters. In particular, $\mathcal{F}$ is never physical because it satisfies the cosmological equations only for a negative w . Point $\mathcal{G}$ represents a physical vacuum solution only for $w > 2/3$ and $w \neq 1$. For these values of w , this point is associated with an expansion whose rate cannot be higher than the radiation dominated GR-FLRW solution. Point $\mathcal{H}$ is associated to a non-vacuum solution which resembles a well known Friedmann-GR solution. As n varies, this solution can represent power law inflation, a decelerated Friedmann solution and a contraction. For $n = 2$, this point is linked to a GR-like state, but since it is unstable for every value of the parameters, we conclude that there is no way for this class of scalar tensor cosmologies to approach a stable non-vacuum GR state. In fact, a quick look to the table above reveals that there is no value of the parameters for which any of the finite fixed points are stable. This means that the only non-vacuum attractors for this class of theories are the asymptotic fixed points and their associated Lemaître solution. Therefore we can conclude that the class of cosmologies we treated are doomed either to approach an effectively vacuum state (probably corresponding to thermal death) that corresponds to one of the vacuum attractors or to recollapses towards a Big Crunch.

The most interesting orbits are the ones that “travel” between point $\mathcal{E}$ and $\mathcal{H}$. This is because they could represent cosmic histories in which a Friedmann-like cosmology enters naturally in a phase of accelerated expansion or cosmic histories in which an unstable inflationary phase is followed by a second inflationary phase. The first type of orbits is interesting because they can in principle help solve the incompatibility between the evolution towards a Dark Energy era and the formation of large scale structure. The second ones are interesting because they potentially unify “dark” scalar fields and the inflaton within a single scheme.

Since the phase space is three dimensional it is not easy to check if such orbits actually exist without the use of numerical techniques. However, our results allow one to give some necessary condition for these orbits to exist and to rule out some of them. For example, it is easy to see that there is no value of the parameters for which $\mathcal{E}$ can represent an unstable Friedmann solution and $\mathcal{H}$ an inflationary phase. On the other

hand, for $4 < n < Q_+$ and $w = 0, 1/3$ we have that $\mathcal{H}$ corresponds to an unstable decelerated expansion, $\mathcal{E}$ correspond to a stable power law inflation and these points are not separated by any invariant submanifold. This means that, in principle cosmic histories exist for which a transient Friedmann evolution approaches to a power law inflationary phase in a natural way.

If $4 < n < 6(1+w)/(3w+1)$ and $w = 0, 1/3$ another interesting set of cosmic histories is possible in which we have two inflationary phases, a first one which is unstable associated with $\mathcal{H}$ and a second one which is stable associated with $\mathcal{E}$. Although only a detailed analysis of these cosmic histories can reveal if this last group of cosmic histories also include the deceleration phase necessary for the realization of standard cosmology, this scenario is interesting because it shows that in STG the non-minimally coupled scalar field can act as *both* the inflaton and dark energy. Such a behavior has been also found in a class of higher order gravity models [43] and definitely deserves a more careful investigation. Such a study, will be the topic of a forthcoming paper.

As final comment, it is worth stressing the connection between the model presented above and the String-Dilaton action. Using the transformation (5), which is only a reparametrization for the scalar field, we can pass from the action (1) to (6). This means that all the exact solutions for the scale factor that we have derived together with their stability are also solutions for the String-Dilaton action. Therefore, our analysis also allows us to give details of the dynamics of low-energy string cosmology which could be of great interest in the quest for finding observational constraints for this theory.

Acknowledgments

This research was supported by the National Research Foundation (South Africa) and the Italian *Ministero Degli Affari Esteri-DG per la Promozione e Cooperazione Culturale* under the joint Italy/ South Africa Science and Technology agreement. The authors wish to thank A Starobinsky and G Venturi for their useful comments.

References

- [1] Weinberg S 1972 *Gravitation and Cosmology* (New York: Wiley)
- [2] Fujii Y and Maeda K 2003 *The Scalar-Tensor Theory of Gravity* (Cambridge: Cambridge University Press)
- [3] Jordan P 1955 *Schwerkraft und Weltall* (Braunschwig: Frierich Vieweg und sohn)
- [4] Brans C and Dicke R H 1961 *Phys. Rev.* **124** 925
- [5] S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, *Phys. Lett. B* **639** (2006) 135 [arXiv:astro-ph/0604431]; S. Capozziello, S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **634** (2006) 93 [arXiv:hep-th/0512118].
- [6] Buchbinder I L, Odintsov S D, and Shapiro I L 1992 *Effective Action in Quantum Gravity* (Bristol: IOP Publishing)
- [7] Guth A 1981 *Phys. Rev. D* **23** 347
- [8] La D and Steinhardt P J 1989 *Phys. Rev. Lett.* **62** 376
- [9] La D, Steinhardt P J and Bertschinger E 1989 *Phys. Lett.* **B231** 231
- [10] Peebles P J E and Rathra B 2003 *Rev. Mod. Phys.* **75** 559
- [11] Cardone V F, Troisi A, and Capozziello S 2005 *Phys. Rev. D* **72** 043501 and references therein
- [12] Capozziello S and de Ritis R 1993 *Int. Jou. Mod. Phys. D* **2** 367
- [13] Capozziello S et al 1996 *La Riv. del N. Cim.* **4** 1
- [14] Capozziello S and de Ritis R 1994 *Class. Quantum Grav.* **11** 107
- [15] For a review on the dynamical system theory a useful book is G Sansone and R Conti 1956 “Equazioni Differenziali Non Lineari” (Edizioni Cremonese, Roma) ; for the application of

- dynamical system to cosmology see Wainwright J and Ellis G F R (ed) 1997 *Dynamical systems in cosmology* (Cambridge: Cambridge University Press) and references therein
- [16] Qualitative analysis of cosmological models in Brans-Dicke theory, C.Romero, H.P. Oliveira and J.R.T. Mello Neto, *Astrophysics and Space Science* **158**, 229 (1989)
 - [17] Kolitch S J and Eardley D M 1995 *Ann. Phys. (NY)* **241** 128
 - [18] Kolitch S J 1996 *Ann. Phys. (NY)* **246** 121
 - [19] Santos C and Gregory R 1997 *Ann. Phys. (NY)* **258** 111
 - [20] Holden D J and Wands D 1998 *Class. Quantum Grav.* **15** 3271
 - [21] Amendola L, Litterio M and Occhionero F 1990 *Int. Journ. Mod. Phys. A* **5** 3861
 - [22] D. J. Holden and D. Wands, *Phys. Rev. D* **61** (2000) 043506 [arXiv:gr-qc/9908026].
 - [23] L. Amendola, *Phys. Rev. D* **60** (1999) 043501 [arXiv:astro-ph/9904120].
 - [24] J. D. Barrow and K. i. Maeda, *Nucl. Phys. B* **341** (1990) 294.
 - [25] R. Gannouji, D. Polarski, A. Ranquet and A. A. Starobinsky, *JCAP* **0609** (2006) 016 [arXiv:astro-ph/0606287].
 - [26] Foster S 1998 *Class. Quantum Grav.* **15** 3485
 - [27] Billyard A, Coley A A, Ibanez J 1999 *Phys. Rev.* **D59** 023507
 - [28] Gunzig E *et al.* 2000 *Class. Quantum Grav.* **17** 1783
 - [29] Gunzig E *et al.* 2001 *Phys. Rev.* **D63** 067301
 - [30] Saa A *et al.* 2001 *Int. Journ. Theor. Phys.* **40** 2295
 - [31] Carloni S, Dunsby P K S, Capozziello S and Troisi A 2005 *Class. Quantum Grav.* **22** 4839
 - [32] Leach J A, Carloni S and Dunsby P K S 2006 *Class. Quantum Grav.* **23** 4915
 - [33] Järv L, Kuusk P, M Saal 2007 *Phys. Rev.* **D75** 023505
 - [34] Faraoni V 2005 *Ann. Phys. (NY)* **317** 366
 - [35] Capozziello S, de Ritis R, and Rubano C 1993 *Phys. Lett. A* **177** 8
 - [36] Birrell N D and Davies P C W 1982 *Quantum Fields in Curved Space* (Cambridge: Cambridge University Press)
 - [37] Capozziello S, de Ritis R, Marino A A 1997 *Class. Quantum Grav.* **14** 3243
 - [38] Leach J A, PhD thesis 2008
 - [39] Hartman P 1964 *Ordinary Differential Equations* (New York: Wiley)
 - [40] J. Garcia-Bellido and M. Quiros, *Phys. Lett. B* **243** (1990) 45.
 - [41] T. Damour and K. Nordtvedt, *Phys. Rev. D* **48** (1993) 3436.
 - [42] Capozziello S, Occhionero F and Amendola L 1993 *Int. Journ. Mod. Phys. D* **1** 615
 - [43] M. Abdelwahab, S. Carloni and P. K. S. Dunsby, arXiv:0706.1375 [gr-qc].